ThyroMend™

Hormone Specific Formulation™

DESCRIPTION

ThyroMend™, provided by Douglas Laboratories®, is a formula of iodine containing seaweeds and herbs which contain phytothyroidogenic properties, as well as phyto-thyroid-receptor agonists. This formulation is designed to promote optimal function of thyroid hormones by maintaining the health of thyroid hormone producing tissues and by supporting the healthy function of tissues that respond to thyroid hormones.†

ThyroMend™ is a *Hormone Specific Formulation*™ formulated by Dr Joseph J Collins, created to support the optimal function of specific hormones through the use of hormone specific adaptogens, hormone specific agonists and hormone specific functional mimetics. This formulation may be used to as part of a hormone health program with dietary and nutrient support. In addition, this formulation may be used by clinicians as an adjuvant to support optimal hormone health in patients who have been prescribed bioidentical hormone replacement therapies, including thyroid replacement therapy.

FUNCTIONS

The primary functions of ThyroMend™ are to support the natural production of thyroid hormones by the thyroid gland, and to support how tissues throughout the body respond to thyroid hormones. This is accomplished by providing support for each of the seven steps involved in thyroid function.
The synergistic combination of specific seaweeds and herbs in ThyroMend™ support these *Seven Important Actions* associated with optimal thyroid health:

1) Provide bio-available iodine for thyrocytes†
2) Promote iodine uptake by thyrocytes through interactions with sodium-iodide-symporter (NIS) proteins †
3) Promote T3 & T4 production and secretion from thyrocytes †
4) Promote conversion of thyroxine (T4) to the more bioactive triiodothyronine (T3) by liver cell and other peripheral cells †
5) Promote RXR/TR heterodimerization of thyroid receptors on target cells †
6) Promote binding of thyroid hormone receptors to DNA by RXR/TR heterodimers †
7) Promote action T3 on target gene expression. †

By supporting thyroid functions, this synergistic formulation supports the health of tissues which respond to optimal thyroid health. This support can also help maintain ideal metabolic rate, maintain normal blood lipids and sugar levels, and support memory and mood. †

INDICATIONS

ThyroMend™ may be a useful dietary supplement for individuals wishing to support healthy thyroid production and function.

FORMULA (#201381)

Serving size…….2 v.caps…….4 v.caps
Iodine.....200 mcg…….. 400 mcg
(from seaweed blend)
Proprietary Blend.....1,300 mg……..2,600 mg
Seaweed Blend (*Ascophyllum nodosum & Fucus vesiculosus,* entire plant), Standardized Guggulipid (*Commiphora mukul,* gum resin, 2.5% guggulsterones), Standardized Bacopa (*Bacopa monnieri,* aerial parts, 20% bacosides), Standardized Ashwagandha (*Withania somnifera,* root, 1.5% withanolides & 1% alkaloids), Standardized Hops*(Humulus lupulus* aerial parts, 5% alpha bitter acid), *Coleus forskohlii* (root, standardized to 10% forskolin), Sage (*Salvia officinalis,* leaf), & Standardized Rosemary extract (*Rosmarinus officinalis,* aerial parts, 6% carnosic acid)

Other Ingredients: Hydroxypropyl methylcellulose (capsule), dicalcium phosphate, vegetable stearate, silica

ThyroMend™

Hormone Specific Formulation™

SUGGESTED USE

As a dietary supplement, adults may take ***2*** capsules each day with food for **1 to 2 weeks** or as directed by your healthcare professional. The dose may then be increased to ***4*** capsules each day with food for **2 to 4 months** or as directed by your healthcare professional. After **2 to 4 months** dosage may be lowered back down to ***2*** capsules each day with food and may continue on that dosage as needed or as directed by your healthcare professional.

SIDE EFFECTS

WARNING: Keep out of reach of children. Not to be used by pregnant or nursing women without consent of their healthcare professional.

STORAGE

For optimal storage conditions, store in a cool, dry place.
(59° – 77 F / 15 -25 C)

REFERENCES

Caldwell KL, Jones R, Hollowell JG. Urinary iodine concentration: United States National Health And Nutrition Examination Survey 2001-2002. Thyroid. 2005 Jul;15(7):692-9.
Borak J. Adequacy of iodine nutrition in the United States. Conn Med. 2005 Feb;69(2):73-7.
Gaitan JE, Mayoral LG, Gaitan E. Defective thyroidal iodine concentration in protein-calorie malnutrition. J Clin Endocrinol Metab. 1983 Aug;57(2):327-33.
Campbell WR. Diet and nutrition: iodine in normal nutrition. Can Med Assoc J. 1939 Jan; 40(1): 77-79.
Butler MR. Comparison of the chemical composition of some marine algae. Plant Physiol.
1931 Apr; 6(2): 295-305.
Panda S, Kar A. Gugulu (Commiphora mukul) induces triiodothyronine production: possible involvement of lipid peroxidation. Life Sci. 1999;65(12):PL137-41.
Panda S, Kar A. Guggulu (Commiphora mukul) potentially ameliorates hypothyroidism in female mice. Phytother Res. 2005 Jan;19(1):78-80.
Brobst DE, Ding X, Creech KL, Goodwin B, Kelley B, Staudinger JL. Guggulsterone activates multiple nuclear receptors and induces CYP3A gene expression through the pregnane X receptor. J Pharmacol Exp Ther. 2004 Aug;310(2):528-35. Epub 2004 Apr 1.
Lee S, Privalsky ML. Heterodimers of retinoic acid receptors and thyroid hormone receptors display unique combinatorial regulatory properties. Mol Endocrinol. 2005
Steiner M, Priel I, Giat J, Levy J, Sharoni Y, Danilenko M. Carnosic acid inhibits proliferation and augments differentiation of human leukemic cells induced by 1,25-dihydroxyvitamin D3 and retinoic acid. Nutr Cancer. 2001;41(1-2):135-44.
Danilenko M, Wang X, Studzinski GP. Carnosic acid and promotion of monocytic differentiation of HL60-G cells initiated by other agents. J Natl Cancer Inst. 2001 Aug 15;93(16):1224-33.
Rutherford DM, Nielsen MP, Hansen SK, Witt MR, Bergendorff O, Sterner O. Isolation and identification from Salvia officinalis of two diterpenes which inhibit t-butylbicyclophosphoro[35S]thionate binding to chloride channel of rat cerebrocortical membranes in vitro. Neurosci Lett. 1992 Feb 3;135(2):224-6.
Panda S, Kar A. Changes in thyroid hormone concentrations after administration of ashwagandha root extract to adult male mice. J Pharm Pharmacol. 1998 Sep;50(9):1065-8.
Laurberg P. Forskolin stimulation of thyroid secretion of T4 and T3. FEBS Lett. 1984 May 21;170(2):273-6.
Kar A, Panda S, Bharti S. Relative efficacy of three medicinal plant extracts in the alteration of thyroid hormone concentrations in male mice. J Ethnopharmacol. 2002 Jul;81(2):281-5.
Radovic B, Schmutzler C, Kohrle J. Xanthohumol stimulates iodide uptake in rat thyroid-derived FRTL-5 cells. Mol Nutr Food Res. 2005 Sep;49(9):832-6.

ThyroMend™

Hormone Specific Formulation™

For more information on ThyroMend™ visit **Douglaslabs.com** or **ThyroMend.com**

Hormone Specific Formulation™ is a trademark of Your Hormones, Inc.

† These statements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by
Douglas Laboratories
600 Boyce Road
Pittsburgh, PA 15205
800-245-4440
douglaslabs.com

You trust Douglas Laboratories.
Your patients trust you.

© 2012 Douglas Laboratories. All Rights Reserved